

A Efficient Synthesis of Nanoparticle of Some Benzothiazoles Derivative of Glucose


K. M. Kalange, Ashish G. Sarap, P.T. Agrawal

Department of Chemistry,
Shri R. L. T. College of Science, Akola-444001
(Maharashtra) India

Abstract :-

Benzothiazoles have played an important role in the field of biochemistry and medicinal chemistry due to their highly pharmaceutical and biological activity. The development of synthetic processes is undoubtedly one of the most significant problems facing researchers. Nanotechnology also involves the synthesis of nanoparticles. These compounds arouse interest as potential biologically active substances and versatile intermediates for preparing various derivatives. To achieve the principle of green chemistry process, it leads to in search of green synthesis of nanoparticles. Here we have synthesized 1-Tetra-O-acetyl- β -D-glucosyl-3-(2)- benzothiazolyl thiocarbamides by reaction of Tetra-O-Acetyl-B-D-glucosyl isothiocyanate with various substituted benzothiazole. The identities of newly synthesis compounds have been established based on usual chemical transformation and U.V, IR, NMR, Mass and Partical Size analysis Analytical studies.
Keywords: TAG Isothiocyanate , substituted benzothiazole and 1-Tetra-O-acetyl- β -D-glucosyl-3-(2)- benzothiazolyl thiocarbamides nanoparticles.

Graphical Abstract :-




Introduction :-

Benzothiazole is a representative class of sulfur-containing heterocycles and involves a benzene ring fused to a thiazole ring. The benzothiazole ring system was originally found in various marine and terrestrial natural compounds, which is widely used as vulcanization accelerators, antioxidants, plant growth regulators, anti-inflammatory agents, enzyme inhibitors, imaging reagents, fluorescence materials, and electroluminescent devices due to its highly pharmaceutical and biological activity [1,2,3,4]. Especially, benzothiazole plays an important role in the field of medicinal chemistry and renders an extensive range of biological activities including anti-cancer [5,6], anti-bacterial [7,8], anti-tuberculosis [9,10], anti-diabetic [11], anthelmintic [12], anti-tumor [13,14,15], anti-viral [16,17], anti-oxidant [18], anti-inflammatory [19,20], anti-glutamate and anti-parkinsonism [21], anticonvulsant [22], muscle relaxant activities [23], neuroprotective [24], inhibitors of several enzymes and so on [25]. Hence, the synthesis of benzothiazoles is of considerable interest due to their potent and significant biological activities and great pharmaceutical value.

Nanotechnology, as defined by size, is naturally very broad, including the field of science as diverse as surface science, organic chemistry, molecular biology, semiconductor physics, energy storage, microfabrication, molecular engineering, etc. The associated research and applications are equally diverse, ranging from extensions of conventional device physics to completely new approaches based upon molecular self-assembly, from developing new materials with dimensions on the nanoscale to direct control of matter on the atomic scale. Nanotechnology may create many new materials and devices with various applications, such as in Nanomedicines, Nanoelectronics, and biomaterial energy production and consumer products.

Results And Discussion:**Nanoparticles:**

A nanoparticle is a sub-classification of the ultrafine particle with lengths in two or three dimensions greater than 0.001 micrometer (1 nanometer) and smaller than about 0.1 micrometer (100 nanometers) and which may or may not exhibit a size-related intensive property. This term is a subject of controversy regarding the size range and the presence of a size-related property. Current usage emphasizes size and not properties in the definition. The length scale may be a hydrodynamic diameter or a geometric length appropriate to the intended use of the nanoparticle. The chemistry of thiourea of carbohydrates is extensively elaborated and well documented. These compounds arouse interest as potential biologically active substances and versatile intermediates for preparing various derivatives

Reaction Scheme:-

Here Tetra-O-Acetyl-β-D-Glucosyl-3-substitute benzothiazolyl thiocarbamide was synthesized by both methods; conventional and Microwave method. Tetra-O-acetyl-β-D-glucosyl isothiocyanate was react with Substitute benzothiazole in benzene medium after that reaction mixture was titurate several times with petroleum ether. The product is confirmed based on the melting point and other studies.

Experimental

Melting points were recorded on electro thermal melting point apparatus are uncorrected. Specific rotations were measured on Equip-Tronic digital polarimeter model no. EQ 800 at 30°C in CHCl₃. IR spectra were recorded on a Perkin Elmer spectrometer. ¹H NMR were obtained on a Bruker DRX-300 (300 MHz FT NMR) NMR spectrometer in CDCl₃ solution with TMS as an internal reference. The mass spectra were recorded on a DART mass spectrometer were recorded. Partical Size was analysis by Malvern particalsize analyzer.

I) Preparation of Tetra-O-acetyl-β-D-glucopyranosyl isothiocyanate :

This has been prepared by the interaction of tetra-O-acetyl-α-D-glucopyranosyl bromide and lead thiocyanate, the former was prepared according to the procedure described earlier. Details of typical experimental are as follows :

a) Microwave assisted preparation of glucose penta acetate :-

Peracetylation of glucose to give the acetyl derivative with small excess of acetic anhydride under the catalyst of either Potassium or Sodium acetate (anhydrous) was found practically quantitative in less than 15 min with microwave heating.

Herein, we reported first time peracetylation of glucose in molecular proportion of acetic anhydride (30 ml) using catalyst sodium acetate 0.8 gm. Under Microwave heating the reaction was complete less than 10 min. Product was isolated by pour in ice cold water with constant stirring and cooling.

The glucose penta acetate is separated out; purification of product was done under water, ethanol system. Melting point of Glucose penta acetate was found to be 110°C.

b) Synthesis of Tetra-O-acetyl- α -D-glucopyranosyl bromide :

The finely powdered glucose pentaacetate (21.0g) was added gradually to the brominating reagent. After the addition the flask was kept for 2 hr. at room temperature. The reaction mixture was mixed with chloroform (50 ml) then the mixture was shaken vigorously for about 15 min. The resultant mixture was pour in ice cold water.

The chloroform layer was then separated. It was washed several time with aqueous sodium bicarbonate to removed excess of acetic acid followed by the aqueous sodium meta-bisulphate to remove the excess of bromine and finely 2-3 times with water. To the chloroform layer addition of petroleum ether afforded a solid (15 g). This solid was expected tetra-O-acetyl- α -D-glucopyranosyl bromide, it was crystallized from ethanol, m.p. 88-90°C.

d) Preparation of lead thiocyanate :

Lead thiocyanate was prepared by mixing aqueous solution of lead nitrate and ammonium thiocyanate. The white granular lead cyanate was filtered washed with distilled water and dried at 50°C.

Preparation of Tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate :

To a suspension of tetra-O-acetyl- α -D-glucopyranosyl bromide (21 g) in sodium dried xylene (80 ml) was added lead cyanate (15g). The reaction mixture was refluxed gently for 3 hr. with frequent shaking. This solution was then cooled and liberated lead bromide was removed by filtration. The xylene filtrate was then treated with petroleum ether (60-80°) with stirring, a pale yellow solid obtained (12 g). This solid was expected tetra-O-acetyl- β -D-glucopyranosyl isocyanate. It was purified by dissolving it in a minimum quantity of chloroform and reprecipitating with petroleum ether. m.p 115-120°C

Table No-2:- Study of synthesis of 1-tetra-O-acetyl- β -D-Glucosyl isothiocyanate under microwave irradiation¹¹

Sr. No.	Amount of G-Bromide	Amount of Xylene	Time	Power in watt	Temp. °C	% Yield
1	10.0 gm	80 ml	35 min	P-70	120-130	90%
2	20.0 gm	120 ml	40 min	P-80	130-140	80%
3	30.0 gm	150 ml	40 min	P-80	135-145	65%

- Lead thiocyanate was taken in equimolar proportion of G-Bromide.
- G-Bromide :- 1-tetra-O-acetyl- β -D-Glucosyl-Bromide

Experiment No. 1:- Tetra-O-acetyl- β -D-glucosyl-(3)-2-substituted benzothiazolylthiocarbamides

A benzene solution of tetra-O-acetyl- β -D-glucosyl isothiocyanate (0.005 M, 1.9 g, in 20 ml) was mixed with the suspension of 2-amino benzothiazole (0.005 M, 0.75 g in 10 ml) and mixture was under microwave irradiation After, benzene was distilled off and sticky mass obtained as a residue was triturated several time with petroleum ether to afford white solid. It was crystallized from ethanol water, m.p. 135-142°C. [Found: C, 52.00; H, 4.85; N, 8.14; S, 6.23, C₂₂H₂₅O₁₀N₃S. requires; C, 52.07, H, 4.93; N, 8.28; S, 6.31%].

The product was found soluble in ethanol, acetone, chloroform and benzene while insoluble in water and petroleum ether. It charred when heated with conc. sulphuric acid. It was non-desulphurisable when boiled with alkaline plumbite solution. It was found to be optically active and its specific rotation $[\alpha]_D^{29} = +74.46^\circ$ (c, 0.94 in chloroform). The purity was checked by TLC and recorded Rf value 0.86 (CCl₄ : EtOAc, 3:2.1).

Analytical And Spectral Data Of Compounds:**Synthesis of 1-Tetra-O-acetyl- β -D-glucosyl-3-(2)- amino benzothiazolyl thiocarbamides (IIIa).**

(IIIa) IR (KBr) : ν 3374 (N-H), 2969 (Ar-H) 1743 (C=O), 1588 (C=N), 1382 (C-N), 1239 (C-O) and 753 cm⁻¹ (C-S); 1H NMR (CDCl₃) : δ 7.57-7.20 (m, 5H Ar-H), 6.3-6.2 (s 1H N-H), 5.1-5.5 (m, 7H Glucopyranosyl ring), 2.1-1.9 (m, 12 H, 4COCH₃); Mass : m/z 508 (M⁺), 331, 169, 109, Anal. Calcd. for C₂₂H₂₅N₃O₈S₂; C, 48.97; H, 4.63; N, 8.24; S, 6.31; found C, 48.84; H, 4.59; N, 8.45; S 6.51%

Synthesis of 1-tetra-O-acetyl- β -D-glucosyl-3(2)-4-chloro benzothiazolyl thiocarbamide (IIIb).

(IIIb) IR (KBr) : ν 3340 (N-H), 2966 (Ar-H), 1745 (C=O), 1615 (C=N), 1378 (CN), 1231 (C-O) and 753 cm⁻¹ (C-S) ; 1H NMR (CDCl₃) : δ 7.7-7.6 (m, 1H, N-H), 7.4 - 7.1 (m, 4H, Ar-H), 5.1-4.7 (m, 7H, glucopyranosyl ring), 2.2-1.9 (m, 12H, 4 COCH₃); Mass : m/z 544 (M⁺), 331, 169, 109; Anal. Calcd., C₂₂H₂₄N₃O₈S₂-Cl, C, 46.07; H, 4.18; N, 7.72; S, 5.88 Found C, 46.13; H, 4.23; N, 7.21; S, 5.98%

Synthesis of 1-Tetra-O-acetyl- β -D-glucosyl-3-(2)-4-methyl benzothiazolyl thiocarbamide (IIIe) :

(IIIe) IR (KBr) : ν 3468 (N-H), 2997 (Ar-H), 1743 (C=O), 1610 (C=N), 1383 (C-N), 1238 (C-O) and 845 cm⁻¹ (Glucopyranosyl ring); 1H NMR (CDCl₃) : δ 7.4-7.0 (m, 4H, ArH), 5.15 (s, 1H, N-H), 2.3 (m, 3H, Ar-CH₃)

5.4-5.5 (m, 7H, Glucopyranosyl ring), 2.2-1.8 (m, 12H, 4COCH₃) Mass: m/z (Not predictable) 331, 109; Anal. Calcd. for C₂₃H₂₇N₃O₈S₂; C, 49.90; H, 4.33; N, 8.04; S, 6.13; Found C, 49.97; H, 4.26; N, 8.22; S, 6.21%.

Preparation of Nanoparticles of 1-tetra-O-acetyl-β-D-glucosyl-3-(2)-methyl benzothiazolyl thiocarbamide:

Take about 1 gm of 1-tetra-O-acetyl-β-D-glucosyl-3-(2)-methyl benzothiazolyl thiocarbamide and dissolve it completely in the 20ml of solvent in a 250 ml beaker and add poly vinyl alcohol as a stabilizer 1.5ml. Now put this beaker in a sonicator. The highly penetrating acoustic waves are passed through the mixture, which creates high-pressure bubbles in the beaker due to which breakdown of the bulk material took place and desired sized nanoparticles are formed. Then stirred mixture about 6hr. in magnetic stirrer at room temperature. The size determination of nanoparticles is done by the particle size analyzer studies

Characterization of Nanoparticles:

1. Characterization using UV-Spectrophotometer: Single Beam UV-Spectrophotometer with software BI/CI/SP/SB-S-03 of Bio Era make. The UV-Visible Spectroscopy reveals the formation of Nanoparticles. Characterization of Nanoparticles was done using a visible Spectrophotometer by using a model by showing different absorption those from bulk material.

2. Size determination of Glucose Penta Acetate, Nanoparticles by X-ray Diffraction studies (Particle Size Analysis): From the X-ray diffraction, it comes to know that size of nano Glucose Penta Acetate is 97.57 nm.




Table No. 2:- Tetra-O-acetyl-β-D-glucosyl-(3)-2-substituted benzothiazolylthiocarbamides

Sr.No.	-Nano of synthesis of Tetra-O-acetyl-β-D-glucosyl-(3)-2-substituted benzothiazolyl thiocarbamides	M.P. °C	Particle Size
1	1-Tetra-O-acetyl- β -D-glucosyl-3-(2)- amino benzothiazolyl thiocarbamides	135-142	122.2nm
2	Synthesis of 1-tetra-O-acetyl- β -D-glucosyl-3(2)-4-chloro benzothiazolyl thio carbamide	112-120	153nm
3.	Synthesis of 1-Tetra-O-acetyl- β -D-glucosyl-3-(2)-4-methyl benzothiazolyl thio carbamide	152-155	97.57nm

References

- Carroll, A.R.; Scheuer, P.J. pentacyclic alkaloids from a tunicate and its prosobranch mollusk predator *Chelynotussemperi*. *J. Org. Chem.* **1990**, *55*, 4426–4431.
- Gunawardana, G.P.; Kohmoto, S.; Gunasekera, S.P.; McConnel, O.J.; Koehn, F.E. Dercitine, a new biologically active acridine alkaloid from a deep water marine sponge, *Dercitus* sp. *J. Am. Chem.* **1988**, *110*, 4856–4858.
- Noel, S.; Cadet, S.; Gras, E.; Hureau, C. The benzazole scaffold: A SWAT to combat Alzheimer's disease. *Chem. Soc. Rev.* **2013**, *42*, 7747–7762.
- Prajapati, N.P.; Vekariya, R.H.; Borad, M.A.; Patel, H.D. Recent advances in the synthesis of 2-substituted benzothiazoles: A review. *RSC Adv.* **2014**, *4*, 60176–60208.
- Kok, S.H.L.; Gambari, R.; Chui, C.H.; Yuen, M.C.W.; Lin, E.; Wong, R.S.M.; Lau, F.Y.; Cheng, G.Y.M.; Lam, W.S.; Chan, S.H.; et al. Synthesis and anti-cancer activity of benzothiazole containing phthalimide on human carcinoma cell lines. *Bioorg. Med. Chem.* **2008**, *16*, 3626–3631.

6. Heo, Y.; Song, Y.S.; Kim, B.T.; Heo, J.N. A highly regioselective synthesis of 2-aryl-6-chlorobenzothiazoles employing microwave-promoted Suzuki–Miyaura coupling reaction. *Tetrahedron. Lett.* **2006**, *47*, 3091–3094.
7. Alaimo, R.J.; Pelosi, S.S.; Freedman, R. Synthesis and Antibacterial Evaluation of 2-(Substituted Phenylureido)-4-thiocyanatobenzothiazoles. *J. Pharm. Sci.* **1978**, *67*, 281–282.
8. Singh, M.; Singh, S.K.; Gangwar, M.; Nath, G.; Singh, S.K. Design, synthesis and mode of action of some benzothiazole derivatives bearing an amide moiety as antibacterial agents. *RSC Adv.* **2014**, *4*, 19013–19023
9. Čaleta, I.; Grdiša, M.; Mrvoš-Sermek, D.; Cetina, M.; Tralić-Kulenović, V.; Pavelić, K.; Karminski-Zamola, K. Synthesis, crystal structure and antiproliferative evaluation of some new substituted benzothiazoles and styrylbenzothiazoles. *Il Farmaco.* **2004**, *59*, 297–305.
10. Das, J.; Moquin, R.V.; Lin, J.; Liu, C.; Doweiko, A.M.; Defex, H.F.; Fang, Q.; Pang, S.; Pitt, S.; Shen, D.R.; et al. Discovery of 2-amino-heteroaryl-benzothiazole-6-anilides as potent p56lck inhibitors. *Bioorg. Med. Chem.* **2003**, *13*, 2587–2590.
11. Su, X.; Vicker, N.; Ganeshpillai, D.; Smith, A.; Purohit, A.; Reed, M.J.; Potter, B.V.L. Benzothiazole derivatives as novel inhibitors of human 11 β -hydroxysteroid dehydrogenase type 1. *Mol. Cell. Endocrinol.* **2006**, *248*, 214–217.
12. Sreenivasa, G.M.; Jayachandran, E.; Shivakumar, B.; Jayaraj, K.K.; Vijay, K.M.M.J. Synthesis of bioactive molecule fluoro benzothiazole comprising potent heterocyclic moieties for anthelmintic activity. *Arch. Pharm. Sci. Res.* **2009**, *1*, 150–157.
13. Hutchinson, I.; Jennings, S.A.; Vishnuvajjala, B.R.; Westwell, A.D.; Stevens, M.F.G. Antitumor benzothiazoles.
14. 16. Synthesis and pharmaceutical properties of antitumor 2-(4-aminophenyl) benzothiazole amino acid prodrugs. *J. Med. Chem.* **2002**, *45*, 744–747.
15. Bradshaw, T.D.; Westwell, A.D. The development of the antitumour benzothiazole prodrug, Phortress, as a clinical candidate. *Curr. Med. Chem.* **2004**, *11*, 1009–1021. [
16. Yoshida, M.; Hayakawa, I.; Hayashi, N.; Agatsuma, T.; Oda, Y.; Tanzawa, F.; Iwasaki, S.; Koyama, K.; Furukawa, H.; Kurakata, S.; et al. Synthesis and biological evaluation of benzothiazole derivatives as potent antitumor agents. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 3328–3332.
17. Vicini, P.; Geronikaki, A.; Incerti, M.; Busonera, B.; Poni, G.; Cabras, C.A.; Colla, P.L. Synthesis and biological evaluation of benzo [d] isothiazole, benzothiazole and thiazole Schiff bases. *Bioorg. Med. Chem.*, **2003**, *11*, 4785–4789.
18. Nagarajan, S.R.; Crescenzo, G.A.; Getman, D.P.; Lu, H.F.; Sikorski, J.A.; Walker, J.L.; McDonald, J.J.; Houseman, K.A.; Kocan, G.P.; Kishore, N.; et al. Discovery of novel benzothiazole sulfonamides as potent inhibitors of HIV-1 protease. *Bioorg. Med. Chem.* **2003**, *11*, 4769–4777.
19. Cressier, D.; Prouillac, C.; Hernandez, P.; Amourette, C.; Diserbo, M.; Lion, C.; Rima, G. Synthesis, antioxidant properties and radioprotective effects of new benzothiazoles and thiazoles. *Bioorg. Med. Chem.* **2009**, *17*, 5275–5284.
20. Dogruer, D.S.; Ünlü, S.; Şahin, M.F.; Yqilada, E. Anti-nociceptive and anti-inflammatory activity of some (2-benzoxazolone-3-yl and 2-benzothiazolone-3-yl) acetic acid derivatives. *Il Farmaco.* **1998**, *53*, 80–84.
21. El-Sherbeny, M.A. Synthesis of certain pyrimido [2, 1-b] benzo-thiazole and benzothiazolo [2,3-b] quinazoline derivatives for in vitro antitumor and antiviral activities. *Arzneimittelforschung.* **2000**, *50*, 848–853.
22. Jimonet, P.; Audiau, F.; Barreau, M.; Blanchard, J.C.; Boireau, A.; Bour, Y.; Coleno, M.A.; Doble, A.; Doerflinger, G.; Huu, C.D.; et al. Riluzole series. Synthesis and in vivo “antiglutamate” activity of 6-substituted-2-benzothiazolamines and 3-substituted-2-imino-benzothiazolines. *J. Med. Chem.* **1999**, *42*, 2828–2843.
23. Siddiqui, N.; Rana, A.; Khan, S.A.; Haque, S.E.; Alam, M.S.; Ahsan, W.; Ahmed, S. Anticonvulsant and Toxicity Evaluation of Newly Synthesized 1-[2-(3, 4-disubstituted phenyl)-3-chloro-4-oxoazetidin-1-yl]-3-(6-substituted-1, 3-benzothiazol-2-yl) ureas. *Acta Chim. Slov.* **2009**, *56*, 462–469.
24. Rajeeva, B.; Srinivasulu, N.; Shantakumar, S.M. Synthesis and Antimicrobial Activity of Some New 2-Substituted Benzothiazole Derivatives. *J. Chem.* **2009**, *6*, 775–779.
25. Danzeisen, R.; Schwalenstoecker, B.; Gillardon, F.; Buerger, E.; Krzykalla, V.; Klinder, K.; Schild, L.; Hengerer, B.; Ludolph, A.C.; Dorner-Ciossek, C.; et al. Targeted antioxidative and neuroprotective properties of the dopamine agonist pramipexole and its nondopaminergic enantiomer SND919CL2x [(+) 2-amino-4, 5, 6, 7-tetrahydro-6-L-propylamino-benzathiazole dihydrochloride]. *J. Pharmacol. Exp. Ther.* **2006**, *316*, 189–199.