

International Journal Of Current Engineering And Scientific Research (IJCESR)

ISSN PRINT-2393-8374,ISSN ONLINE-2394-0697

DOI-10.21276/Ijcesr

Home Editors Recent issues Topic Submission Registration Monographs Contact

IMPACT FACTOR

IARC(JCRR)-0.916
GIF-0.695
SIF-0.73
JIF-0.9
ICV 2015- 63.71

Indexed By



VOLUME 6 ISSUE 1 (PART-III) (JANUARY 2019) | PUBLICATION DATE:25TH JANUARY 2019

DOI:10.21276/Ijcesr

1. SYNTHESIS OF GRAPHENE OXIDE-Fe(III) COMPLEX: A NEW MATERIAL FOR CATALYSIS
Authors: Charali C Brtkara, Sanjay R. Thakare, R. G. Atrama, K. G. Rewatkar
2. NOVEL NANOSTRUCTURE ELECTROLYTIC MATERIALS BY X-SIXED AZO DYES FOR FUEL CELL APPLICATION
Authors: M. M. Fukate, A. B. Bodele, G. N. Chaudhan
3. RAMAN SPECTROSCOPY AND THERMAL STUDIES OF RUBBER-NANOCOMPOSITES SYNTHESIZED FROM STYRENE-BUTADIENE RUBBER AND NANO ALUMINUM OXIDE
Authors: Rani V. Mankar, Vasudeo B. Gurnule
4. SYNTHESIS, CHARACTERIZATION AND ANTIMICROBIAL ACTIVITY OF ZINC METAL COMPLEX HAVING ON DONOR LIGANDS
Authors: Apeksha W. Walke, Niven E. Kothale
5. MODIFIED PROTOCOL FOR RAPID IN VITRO DEVELOPMENT CALLUS OF SYZYGIUM CUMINI (L.)
Authors: P.P. More, B. Y. Patil, G. S. Gond, J. A. Kulkarni
6. PHOTOCATALYTIC DEGRADATION OF INDIGO CARMINE DYE ON COMBUSTION SYNTHESIZED MgZrO3 CATALYST UNDER SOLAR LIGHT IRRADIATION
Authors: C. R. Hatwar, P. S. Deshpande, S. Dalare, L. R. Hatwar, C. B. Talwar, R. S. Bhavsar
7. A CRITICAL REVIEW OF THE MEDICINAL PROPERTIES OF SPILANTHES ACMELLA MLUR
Authors: Chandore Hemant D., Jadhav Dnyandea S.
8. RAPID APPROACH FOR VARIETAL IDENTIFICATION AMONG SESAME GENOTYPE
Authors: Triveni Sarita, Burade Nishant, Harode Vijay, Moghe Sandhya
9. MOSSBAUER, XRD AND THERMALLY INDUCED STRUCTURAL STUDIES ON POLYANILINE NANOCOMPOSITE MATERIAL CONTAINING DIFFERENT CONCENTRATION OF MAGNETITE NANOPARTICLE
Authors: A. K. Barve, S. M. Godegane, M. R. Lanjewar, R. B. Lanjewar
10. EXPLORING THE COMPLEX DIELECTRIC PROPERTIES OF ONE POT CHEMICALLY ENGINEERED CONDUCTING POLYIMIDE
Authors: N. S. Wadkar, S. A. Naghuley
11. AN OVERVIEW ON STRUCTURAL AND MAGNETIC PROPERTIES OF CERTAIN SUBSTITUTIONS IN COBALT FERRITE
Authors: A. M. Shahare, M. S. Bisen, A. V. Bagde, D. S. Choudhary
12. PRODUCTION AND CHARACTERIZATION OF MICROWAVE ASSISTED AND CHEMICALLY IMPREGNATED BIOADSORBENT MATERIAL FROM AZADIRACHTA INDICA BARK
Authors: Nandkishor G. Telkapalwara, Vidyadhar M. Shivankar
13. CHARACTERISTIC STUDY OF HEAVY METAL ION BY USING THIN LAYER CHROMATOGRAPHY WITH HUMIC ACID AND L-METHIONINE AS A MOBILE PHASE
Authors: Meghna H. Jurde, W. B. Gurnule
14. STUDY OF THERMODYNAMIC PARAMETERS OF AMINO ACIDS AT 287K & 292K BY VISCOSITY MEASUREMENT
Authors: Chakar Patil, R. R. Tayade
15. SYNTHESIS, SPECTROSCOPIC CHARACTERIZATION AND ANTIMICROBIAL SCREENING OF SOME NEWLY SYNTHESIZED PROPANE 1,3-DIONE (β-DIKETONES) DERIVATIVES
Authors: Rajendra M. Pathade, Pravin S. Borkhe
16. SYNTHESIS AND CHARACTERIZATION OF 5-SUBSTITUTED NAPHTHALENE 1,3,4-THIAZOLE 2 BENZAMIDE WITH THEIR BIOLOGICAL STUDIES
Authors: S. A. Athawale, S. E. Bhandarkar
17. PREPARATION, CHARACTERIZATION OF COCONUT SHELLS ACTIVATED CARBON USING FeCl3 AS ACTIVATING AGENT AND REMOVAL OF TRANSITION METAL IONS
Authors: V. A. Shirbhate, D. P. Gulwade
18. AFM STUDY OF EGG SAC SILK FIBERS OF THE GIANT WOOD SPIDER NERPHILA PILIPES (ARANEAE, ARANEIDAE)
Authors: Sawarkar A. S., Sawarkar S. B.
19. ELECTRICAL CONDUCTANCE PROPERTIES OF COPOLYMER DERIVED FROM 2-HYDROXYACETOPHENONE-MELAMINE-FORMALDEHYDE
Authors: R. M. Gupta, Punam G. Gupta, W. B. Gurnule





PHYTOCHEMICAL SCREENING OF POLLEN OF *CATHERANTHUSROSEUS* PLANT

Sangole AA,

Department of Botany, Shri. R.L.T. College Science, Akola, Maharashtra India

Abstract

The member of Apocynaceae family of *Catheranthusroseus* plant being used in traditional medicine. To study the phytochemical screening of pollen of the plant *Catheranthusroseus* have medicinal properties was carried out. Pollen material was successfully extracted with different organic solvent. The phytochemical screening reveals the presence of alkaloids, steroids, flavinoids, tannins saponins and carbohydrates in them. The result reveals the presence of alkaloids and steroids are reported to have medicinal properties. The presence of these phytochemical can be correlated with the medicinal potential of this plant.

Keywords: Phytochemicals, Pollen, *Catheranthusroseus*.

Introduction

Medicinal plants play a major role in the medicinal needs of about 70% populations in developing countries, which serve as an important resource for the treatment of various maladies and illnesses (Ngari et al., 2010). Globally, about 85% of the traditional medicines used by different ethnic groups inhabiting various terrains for primary healthcare are derived from plants, especially in India; medicinal plants are widely used by all sections of the population with an estimated 7500 species of plants used by several ethnic communities (Farnsworth, 1988). The plant is being used by the local peoples and tribal of Maharashtra as ethno medicine on various ailments. This plant is also being used for its

anti-inflammatory, anti-diarrheal properties by various communities in Indian subcontinent and also across the world. The present study was designed to evaluate the fundamental phytochemical constituents of this wild medicinal plant. It is one of the well-known medicinal plants that have its origin from the continent of Africa. The plant is commonly known as the Madagascar periwinkle or basically *Vincarosea*. It is a tender, perennial plant, which grows as a herb or sub-shrub, growing to approximately 1 metre in height. Leaves simple, petiolate. Flowers are commonly white or pink.

They are known to have various biological activities such as antimicrobial, antifungal, antioxidant, etc. The important bioactive components in plants are usually the secondary metabolites such as alkaloids, flavonoids, tannins and other phenolic compounds (Edeoga et al., 2005). The Medicinal plants have potent phytochemical components which are important source of antibiotic compounds and are responsible for the therapeutic properties (Jeeva et al., 2011; Jeeva and Johnson, 2012; Florence et al., 2012 & 2014; Joselin et al., 2012 & 2013; Sainkhediya and Ray, 2012; Sumathi and Uthayakumari, 2014). As pollen is very important entity carrying paternal genome to next generation, it is quite likely that different important constituents may be present in them. Therefore, preliminary phytochemical screening of pollen grains was carried out in all the plants under investigation.

Table 1: Medicinal uses of the plants in the study

Plant species	Common name in English	Traditional uses
<i>Catheranthusroseus</i>	Rose Periwinkle	Anti diabetic, anti cancer ,leaf decoction use as indigestion, for induce vomiting

Material and Methods:

For phytochemical screening sufficient pollen grains material was collected from *Catheranthusroseus* plants. It was dried, weighed and used for Extraction of constituents.

Extraction of Constituents:

The weighed quantity of pollen material was successfully extracted with different organic solvents. The solvents used for successive extraction were petroleum ether (60 to 80°C) and alcohol. The solvent was removed by Evaporation from the extract and the residue left was weighed from this, the percentage yield to solvent was calculated. Then it was subjected to various chemical tests as follows.

I- Test for Sterols:

(a) Salkowski reaction: - Few mg of residue of each extract was taken in 2ml of chloroform and 2ml of sulphuric acid was added from the side of the test tube. The test tube was shaken for few minutes. The development of red color in the chloroform layers indicated the presence of sterols.

(b) Liebermann-Burchard Reaction: - Few mg of residue was dissolved in chloroform and few drops of acetic anhydride were added followed by conc. sulphuric acid from the side of the test tube. A transient colour development from red to blue and finally green indicated the presence of sterols.

II- Test of Alkaloids:

Few mg of residue of each extract was taken separately in 5ml of 1.5% v/v hydrochloric acid and filtered. These filtrates were then used for testing alkaloids.

a) Mayer's Reagent test: - To a little of the test filtrate taken in a watch glass a few drops of Mayer's reagent was added. Formation of cream coloured ppt. showed the presence of alkaloids.

b) Wagner's Reagent: - few ml of test filtrate was taken in test tube and to this Wagner's reagent was added, a brown flocculent

precipitate was formed indicating the presence of alkaloids.

III Test of Proteins:

a. Bureit test: - A few ml of test residue was taken in water and 1 ml of 4% sodium hydroxide solution was added followed by a 1% solution of copper sulphate. Violet Pink colour development indicates the presence of protein.

b. Xanthoprotein test: - Few 2 ml of extract was taken to this was added 5ml of Nitric acid, white precipitate indicates the presence of protein.

IV Test for Saponins:

a. Foam Test: - A few mg of test residue was taken in a test tube and shaken vigorously with a small amount of sodium bicarbonate and water. If stable, characteristic honey comb like froth is obtained, saponins are present.

V Test for Caumarins:

A small amount of test residue moistened with water was taken in test tube. The mouth of test tube was covered with paper mounted with dilute sodium hydroxide solution. The covered test tube was placed in boiling water bath for several minutes. The paper was removed and exposed to ultraviolet light. If yellowish green fluorescence is obtained, the caumarins are present.

VI Test for Tanins:

The test residue of each extract was taken separately in water, warmed and filtered. Tests are carried out with the filtrate using following reagents.

a) Ferric chloride solution: - Few drops of ferric chloride were added to a little of the above test filtrate. If greenish or bluish black colour is obtained, Tanins are present.

b) Lead acetate Test: - Few drops of lead acetate solution were added to the test filtrate. If precipitate is obtained, tannins are present.

VII Test for Flavinoids:

a) Shinoda Test: - A small quantity of test residue was dissolved in 5ml of ethanol (95%) and added small piece of magnesium ribbon followed by few drops of concentrated hydrochloric acid, colours ranging from orange to red (flavones), red to crimson (flavonols) crimson to magenta (flavonones) and occasionally to green or blue are developed within 3 minutes if flavinoids are present.

VIII Test for sugars:

Fehlings Solution Test: - The test residue solution was heated and drop by drop Fehling solution was added to it and warmed. If red precipitate of cuprous oxide is obtained, reducing sugars including all monosaccharides and many disaccharides are present

Sr. No.	Test	Petroleum Ether
1	Sterols: a) Salkowaski reaction	+
	b) Liebermann-Burchard reaction	+
2	Alkaloids: a) Mayer's reagent	+
	b) Wagner's reagent	+
3	Proteins: a) Bureit test	-
4	Saponins: a) Foam test	-
5	Caumarins	-
6	Tannins: a) Feric Chloride reagent	-
	b) Leads acetate reagent	-
7	Flavinoides: a) Shinodha test	-
8	Carbohydrates: a) Fehling's Test	-

RESULT AND DISCUSSION

Catheranthusroseus pollen were collected, the pollen dried under shaded condition at room temperature. Pollen of *Catheranthusroseus* showed the presence of Steroids and Alkaloids which are reported to have medicinal properties. The earliest chemical investigation of *Catheranthusroseus* was carried out the presence of alkaloidal constituents in them. More than 100 alkaloids have been isolated from the plant and their biological uses have been widely explored. (Trease and Evans, 1978). These alkaloids possess antifibrillic, hypotensive, sedative and tranquilizing properties. The results confirm the presence of constituents which are known to exhibit medicinal as well as physiological activities (Mukeshwar et al 2011). The presence of some

of these compounds has also been confirmed to have antimicrobial activity. Hence it could be inferred that the plant extracts could be a source for the industrial manufacture of drugs useful in the chemotherapy of some microbial infection

Acknowledgments

The authors would like to thank department of Botany shri R.L.T. College of Science, Akola Maharashtra, India for providing facilities for research work

REFERENCES

- Edeoga HO, Okwu DE, Mbaebie BO, Phytochemical Constituents of some Nigerian Medicinal plants, African journal of Biotechnology, 4, 2005, 685-688.

- Farnsworth NR, 1988. Preliminary phytochemical screening and HPLC Analysis of Flavonoid from Methanolic Extract of Leaves of *Annonasquamosa*. Screening plants for new National Academy Press, Washington, DC, Pp 83-97.
- Florence AR, Joselin J and Jeeva S, 2012. Intraspecific variation of bioactive principles in select members of the genus *Clerodendrum* L. *Journal of Chemical and Pharmaceutical Research*, 11: 4908- 4914.
- Florence AR, Joselin J, Brintha TSS, Sukumaran S and Jeeva S, 2014. Preliminary phytochemical studies of select members of the family Annonaceae for bioactive constituents. *Bioscience discovery*, 5(1): 85-96
- Jeeva S, Johnson M, Aparna JS and Irudayaraj V, 2011. Preliminary phytochemical and antibacterial studies on flowers of selected medicinal plants. *International Journal of Medicinal and Aromatic Plants*, 1(2), 107-114.
- Jeeva S and Johnson M, 2012. Antibacterial and phytochemical studies on *Begonia flaccifera* Bedd. flower. *Asian Pacific Journal of Tropical Biomedicine*, 1(S1): S151-S154.
- Joselin J, Brintha TSS, Florence AR and Jeeva S, 2012. Screening of select ornamental flowers of the family Apocyanaceae for phytochemical constituents. *Asian Pacific Journal of Tropical Disease*, 2: 1-
- Joselin J, Brintha TSS, Florence AR and Jeeva S, 2013. Phytochemical evaluation of Bignoniaceae flowers. *Journal of Chemical and Pharmaceutical Research*, 5(4): 106-111.
- Mukeshwar Pandey, Mousumi Debnath and Etal, *Phytomedicine: An ancient approach turning into future potential source of therapeutics*, *Journal of Pharmacognosy and phytotherapy*, 3(1), 113-117 (2011)
- Ngari EW, Chiuri LW, Kariuki ST and Hockett S, 2010. Ethnomedicine of Ogiek of River Njoro Watershed. *Ethnobotany Research and Applications*, 8: 135-152.
- Sainkhediya J and Ray S, 2012. Preliminary study of flowering plant diversity of Nimar region. *Bioscience Discovery*, 3(1): 70-72.
- Sumathi BM and Uthayakumari F, 2014. GC MS analysis of Leaves of *Jatropha maheswarii* Subram & Nayar. *Science Research Reporter*, 4(1): 24-30.
- Trease, G.E. and W.C. Evans, 1978. *Text Book of pharmacognosy*; (11th Edition) P-475.