

THERMODYNAMIC STUDIES ON MOLECULAR INTERACTIONS IN AQUEOUS SOLUTIONS OF BARBITURIC ACID, 1,3-DIMETHYL BARBITURIC ACID AND THIOBARBITURIC ACID

Vinod D. Deotale^a, Madhukar G. Dhonde^{*b}, Satish G. Goswami^b, Santosh D. Desarkar^c, Avinash L. Puyad^c

^aDepartment of Chemistry, Shri Radhakisan Laxminarayan Toshniwal College of Science, Akola, MS, India, 444001

^bDepartment of Chemistry, Shri Mathuradas Mohota Science College, Nagpur, MS, India, 440024

^cSchool of Chemical Sciences, Swami Ramanand Teerth Marathwada University, Nanded 431 606, MS, India

Abstract

The densities, ultrasonic velocities and refractive indices of barbituric acid, 1,3-dimethyl barbituric acid and thiobarbituric acid in aqueous solutions have been measured at 37 °C. The volumetric and acoustical properties were calculated from densities and ultrasonic velocities in order to understand the interactions between barbituric acid-water, 1,3-dimethyl barbituric acid-water and thiobarbituric acid-water. The quantum chemical calculations of barbituric acid, 1,3-dimethyl barbituric acid and thiobarbituric acid in gas phase and in water performed employing GAUSSIAN 09 programme. Energies, bond lengths, IR frequencies of selected interacting groups are reported for studying the solute-solvent interactions.

Keywords:- Thermodynamic properties, Molecular interaction, Compressibility, Barbiturate, DFT

I. INTRODUCTION

Drug-macromolecular interactions are an important phenomenon in physiological media such as blood, membranes, intra and extracellular fluids. The processes of drug transport, protein-binding and anaesthesia are few examples where drug and bio macromolecules appear to interact in an important and vitally significant manner. Thermodynamic properties are generally convenient parameters for interpreting solute-solvent and solute-solute interactions in the solution phase [1-6]. Fundamental properties such as enthalpy, entropy and Gibbs energy represent the macroscopic state of the system and interpretation of these macroscopic properties in terms of molecular phenomena is generally difficult. Sometimes, higher derivatives of these properties can be interpreted more effectively in terms of molecular interactions.

However, some drug effects are non-receptor mediated and are caused by the particular physical or chemical properties of the drug molecule. To firmly grasp the concepts of how desired and deleterious effects are induced in the body by a drug molecule requires an understanding of where and how these molecules interact.

The study of volumetric, acoustical and optical properties of biomolecules in aqueous and aqueous-cosolute solutions provide significant information regarding molecular interactions and hydration behaviour of these molecules. The organic salt like disodium tartarate can change the binding trends and hydration behaviour of biomolecules in solution. The changes in molecular environment and molecular interactions involved are reflected in thermodynamic properties. Interactions between drug and macromolecule are important in biophysical chemistry [7-8]. Drug-electrolyte or drug-active organic molecule interactions are significant for pharmacokinetics and pharmacodynamics. The thermodynamic properties and molecular interactions in aqueous solutions of drug in presence

of electrolytes and other cosolutes have been studied [9-13]. The systems involving different interactions including hydrogen bonding interactions has applications in different fields. The densities, ultrasonic velocities and refractive indices of barbituric acid, 1,3-dimethyl barbituric acid and thiobarbituric acid (0.02-010 mol·kg⁻¹) in aqueous solutions have been measured at 37 oC. The volumetric and acoustical properties were calculated from densities and ultrasonic velocities in order to understand the interactions between barbituric acid-water, 1,3-dimethyl barbituric acid-water and thiobarbituric acid-water. The quantum chemical calculations of barbituric acid (BA), 1,3-dimethyl barbituric acid (1,3-DMBA) and thiobarbituric acid (TBA) in gas phase and in water performed employing GAUSSIAN 09 programme. Energies, bond lengths, IR wave numbers of selected interacting groups are reported for studying the solute-solvent interactions.

II. MATERIALS AND METHODS

2.1. Experimental

Solutions of BA, 1,3-DMBA and TBA in water were prepared in double distilled water using an analytical grade balance, Anamed (Model AA-2200, ±0.0001 g). Densities were measured using graduated pycnometer in triplicate for each solution. The ultrasonic velocities were measured using ultrasonic interferometer (M-F05, Mittal Enterprises) at frequency 2 ± 0.0001 MHz. Temperature of solutions was maintained by electronically controlled thermostatic water bath (±0.1 K). Refractive indices were measured using Cyber LAB-Cyber Abbe Refractometer (*Amkette Analytix*, ±0.0002, 1.3000 to 1.7000). We have used the purified drug molecules directly from different source and detail is as given in Table 1.

TABLE 1. SPECIFICATION OF CHEMICALS.

Name of chemical	Molar mass, g·mol ⁻¹	Source	Formula	Mass fraction purity
Barbituric acid (BA)	128.09	HiMedia	C ₄ H ₄ N ₂ O ₃	≥99.00%
1,3-dimethyl barbituric acid (DMBA)	156.14	Sigma-Aldrich	C ₆ H ₈ N ₂ O ₃	≥99.00%
Thiobarbituric acid (TBA)	144.15	Sigma-Aldrich	C ₄ H ₄ N ₂ O ₂ S	≥99.00%

2.2. Theoretical:

Isentropic compressibility κ_s is calculated from density and ultrasonic velocity [14-15] using equation (1)

$$\kappa = \frac{1}{u\rho} \quad (1)$$

For solvent and solvent mixture, the isentropic compressibility is

$$\kappa_0 = \frac{1}{u_0^2 \rho_0} \quad (2)$$

Specific acoustic impedance (Z) is calculated using equation (3).

$$Z = u\rho \quad (3)$$

Relative association (R_A) is calculated using equation (4).

$$R_A = \frac{\rho}{\rho_0} \times \left(\frac{u_0}{u} \right)^{1/3} \quad (4)$$

Where, ρ_0 is density of solvent and ρ is density of solution ($\text{kg}\cdot\text{m}^{-3}$), u_0 is ultrasonic velocity of solvent and u is ultrasonic velocity of solution ($\text{m}\cdot\text{s}^{-1}$), κ_s and κ_0 are isentropic compressibilities of solution and solvent respectively (Pa^{-1}).

2.3. Computational study:

The quantum chemical calculations are performed employing GAUSSIAN 09 programme [16]. The geometries of BA, 1,3-DMBA and TBA in gas phase and in water optimized using density functional calculations using PCM model at B3LYP/6-31G(d) basis set [17-18]. In order to understand the interactions between BA, 1,3-DMBA and TBA and water, the calculations were performed for BA, 1,3-DMBA and TBA in water in presence of Density Functional Theory (DFT).

III. RESULTS AND DISCUSSION

Measured densities, ultrasonic velocities and refractive indices and calculated isentropic compressibility, specific acoustic impedance and relative association of BA, 1,3-DMBA and TBA at 37 °C are reported in Table 2.

TABLE 2. MEASURED DENSITIES, ULTRASONIC VELOCITIES AND REFRACTIVE INDICES AND CALCULATED ISENTROPIC COMPRESSIBILITY, SPECIFIC ACOUSTIC IMPEDANCE AND RELATIVE ASSOCIATION OF BARBITURIC ACID, 1,3-DIMETHYL BARBITURIC ACID AND THIOBARBITURIC ACID AT 37 °C.

m $\text{mol}\cdot\text{kg}^{-1}$	ρ $\text{kg}\cdot\text{m}^{-3}$	u $\text{m}\cdot\text{s}^{-1}$	n	$\frac{Ks}{\times 10^{-10} \text{ m}^{-2}\cdot\text{N}^{-1}}$	Z $\times 10^6 \text{ kgm}^{-2}\cdot\text{s}^{-1}$	R_A
<i>BA + Water</i>						
0.0000	993.37	1528.90	1.3312	4.3066	1.519	1.0000
0.0202	994.42	1535.22	1.3315	4.2667	1.527	0.9997
0.0404	995.39	1537.41	1.3318	4.2504	1.530	1.0002
0.0607	996.34	1539.27	1.3322	4.2361	1.534	1.0007
0.0810	997.27	1540.46	1.3326	4.2256	1.536	1.0014
0.1015	998.15	1542.24	1.3330	4.2121	1.539	1.0019
<i>1,3-DMBA + Water</i>						
0.0000	993.37	1528.90	1.3312	4.2943	1.521	1.0000
0.0202	994.26	1561.39	1.3315	4.1255	1.552	0.9939
0.0405	995.08	1562.08	1.3318	4.1185	1.554	0.9946
0.0608	995.87	1563.13	1.3320	4.1097	1.557	0.9951
0.0813	996.56	1564.17	1.3323	4.1014	1.559	0.9956
0.1019	997.17	1566.43	1.3325	4.0870	1.562	0.9957
<i>TBA + Water</i>						
0.0000	993.37	1528.90	1.3312	4.3066	1.519	1.0000
0.0201	994.95	1545.04	1.3315	4.2104	1.537	0.9981
0.0403	996.49	1547.13	1.3320	4.1925	1.542	0.9992
0.0606	997.95	1548.17	1.3325	4.1807	1.545	1.0004
0.0809	999.37	1549.91	1.3330	4.1654	1.549	1.0015
0.1013	1000.79	1551.87	1.3335	4.1490	1.553	1.0025

Standard uncertainties, u are $u(T)=0.1 \text{ K}$, $u(p)=\pm 2.0 \text{ kPa}$, $u(\rho)=0.08 \text{ kg}\cdot\text{m}^{-3}$ and $u(m)=0.0005 \text{ mol}\cdot\text{kg}^{-3}$.

The density, ultrasonic velocity and refractive index increases with increase in the concentration of BA, 1,3-DMBA and TBA which is due to the increase in the solute-solvent interactions between BA, 1,3-DMBA and TBA and water. Further, the isentropic compressibility of solution decreases

with increase in the concentration of BA, 1,3-DMBA and TBA which indicates the solution becomes compressible with increase in the concentration of BA, 1,3-DMBA and TBA.

The molecular interactions between BA, 1,3-DMBA and TBA and water like hydrogen bonding interactions, dipole-dipole interactions in solution strengthens with BA, 1,3-DMBA and TBA concentration and solution becomes tight which leads to decrease in the compressibility of solution.

The decrease in K_s with concentration of BA, 1,3-DMBA and TBA indicates that water molecule around polar groups in BA, 1,3-DMBA and TBA are less compressible than bulk [19] due to hydrophilic-hydrophilic interactions [20]. The specific acoustic impedance and relative association support the solute-solvent interactions and strengthening of these interactions with BA, 1,3-DMBA and TBA concentration.

The optimized geometrical structures of BA, 1,3-DMBA and TBA in gas phase and in water at B3LYP/6-31G(d) basis set are shown in Figure 1a-c and the geometrical parameters are reported in Table S1-3. Energies and dipole moments of BA, 1,3-DMBA and TBA in gas phase and in water at B3LYP/6-31G(d) basis set are reported in Table 3.




Fig. 1: Optimized structure of a) barbituric acid, b) 1,3-dimethyl barbituric acid and c) thiobarbituric acid in water using PCM model at DFT b3lyp/6-31g(d) basis set


TABLE 3. ENERGIES AND DIPOLE MOMENTS OF BA, 1,3-DMBA AND TBA IN GAS PHASE AND IN WATER AT B3LYP/6-31G(D) BASIS SET.

System	Energy, kcal/mol	Energy difference	Dipole moment	Dipole moment difference
BA-GP	-307517.5877		0.0410	
BA-W	-307526.6311	-9.0434	0.0062	-0.0348
1,3-DMBA-GP	-356854.2941		0.7542	
1,3-DMBA-W	-356861.4641	-7.1700	0.8144	0.0602
TBA-GP	-510143.9823		5.2408	
TBA-W	-510155.4535	-11.4712	7.0460	1.8052


Foot note- GP=gas phase, W=water, BA=barbituric acid, 1,3-DMBA=1,3-dimethyl barbituric acid, TBA-Thiobarbituric acid.

The energy of BA, 1,3-DMBA and TBA decreased by -9.0434, -7.17 and -11.4712 from gas phase to water and these molecules are stabilized. This stabilization of these molecules in water is due to solvation through solute-solvent interaction. The solute-solvent interactions like hydrogen bonding interactions, dipole-dipole interactions between barbituric acid or 1,3-dimethyl barbituric acid or thiobarbituric acid and water molecule hence stabilize the drug molecules in water has been occur. The change in the dipole moment of the molecules from gas phase to water indicates the solute-


solvent interactions have reported. The theoretical and experimental IR spectra of BA, 1,3-DMBA and TBA are shown in Figure 2-3.




BA in gas phase




TBA in gas phase




1,3-DMBA in gas phase



BA in water




TBA in water



1,3-DMBA in water

Fig. 2: Theoretical IR spectra of BA, 1,3-DMBA and TBA in gas phase, water



The theoretical IR spectra of BA, 1,3-DMBA and TBA in gas phase and in water at B3LYP/6-31G(d) basis set are given in Figure 2. The shift in the wave number values (Table 4) of important vibrations like $>C=O$, $>N-H$ in BA, $>C=O$ in 1,3-DMBA and $-O-H$, $-S-H$ in TBA clearly indicates the interactions between water and BA.

The volumetric, ultrasonic, optical properties of barbituric acid, 1,3-dimethyl barbituric Acid and thiobarbituric acid in water at 37 °C have been studied. The computational and IR spectroscopic study of these molecules have also carried out. The experimental and computational studies confirmed the existence of strong solute-solvent interactions like hydrogen bonding and dipole-dipole interaction between barbituric acid or 1,3-dimethyl barbituric acid or thiobarbituric acid and

water. Further, these interactions strengthen with increase in the concentration of barbituric acid or 1,3-dimethyl barbituric acid or thiobarbituric acid. The stabilization of barbituric acid or 1,3-dimethyl barbituric acid or thiobarbituric acid in water is due to solvation through solute-solvent interaction. The change in the IR frequencies from gas phase to liquid water and from solid to liquid water from computational and experimental IR frequencies of barbituric acid or 1,3-dimethyl barbituric acid or thiobarbituric acid confirmed the hydrogen bonding and dipole-dipole interaction.

TABLE 4. THEORETICALLY CALCULATED WAVE NUMBER OF BA, 1,3-DMBA AND TBA IN GAS PHASE AND IN WATER AT B3LYP/6-31G(D) BASIS SET

Functional group	BA		1,3-DMBA		TBA	
	Gas	water	Gas	Water	Gas	water
>C=O	1839.51	1833.05	1834.21	1815.79	-	-
>N-H	3599.30	3590.29	-	-	-	-
-O-H	-	-	-	-	3749.54	3732.91
-S-H	-	-	-	-	2712.03	2720.23

ACKNOWLEDGEMENTS

Authors are thankful to Dr. J. R. Dontulwar, Principal, Shri M. M. College of science, Nagpur for encouragement and providing necessary facilities.

REFERENCES

1. E. Zorebski **Internal pressure as a function of pressure for alkanols**, *Mole, Quantum Acoustics*, 26, (2005), pp. 317-326.
2. N, Santhi, E. M. Sabarathinam **Molecular interaction studies in binary liquid mixtures from ultrasonic data**, *E. J. Chem.*, 7(2), (2010), pp. 648.
3. B. R. Shinde, K. M. Jadhav **Investigation of physicochemical behaviour of ion-solvent interactions using ultrasonic technique in DMSO drug at different temperatures**, *Pharmacologyonline*, 2, (2010), pp. 533.
4. G. Nath, R. Paikaray **Ultrasonic study of binary mixtures of acetone with chlorobenzene at different frequencies**, *Ind. J. Phys.*, 83(9), (2009), pp. 1309.
5. R. Pilani, K. Jayachitra **Ultrasonic study of ternary electrolytic mixtures at 303K, 308K, and 313K**, *Ind. J. Pure and Appl. Phys.*, 46(4), (2008), pp. 251.
6. R. Kumar, S. Jayakumar, V. Kannappan **Study of molecular interaction in binary liquid mixtures**, *Ind. J. Pure and Appl. Phys.*, 46(3), (2008), pp. 169.
7. A. Pal, S. Soni **Volumetric approach to the interaction of diglycine in aqueous solutions of sulphamethoxazole drug at T= 288.15–308.15 K**, *Fluid Phase Equilib.*, 334, (2012), pp. 144.
8. S. Dhondge, S. Zodape, D. Parwate **Volumetric and viscometric studies of some drugs in aqueous solutions at different temperatures**, *J. Chem. Thermodyn.*, 48, (2012), pp. 207.
9. R. T. Sawale, T. M. Kalyankar, R. George, S. D. Deosarkar **Molar refraction and polarizability of antiemetic drug 4-amino-5-chloro-N-(2-(diethylamino)ethyl)-2-methoxybenzamide hydrochloride monohydrate in {aqueous-sodium or lithium chloride} solutions at 30°C**, *J. App. Pharm Sci.*, 6, (2016), pp. 120.
10. A. Pal, S. Soni **Interaction of aqueous solution of sulphamethoxazole drug with Diglycine by volumetric method at different temperatures**, *Phys. Chem.*, 229 (3), (2015), pp. 443.
11. S. Ryshetti, B. K. Chennuri, R. Noothi, S. J. Tangeda, R. L. Gardas **Volumetric properties of betaine hydrochloride drug in aqueous NaCl and KCl solutions at different temperatures**, *Thermochim. Acta*, 597, (2014), pp. 71.
12. S. P. Jengathe, S. S. Dhondge, L. J. Paliwal, V. M. Tangde, S. Mondal **Effect of sodium chloride and myo-inositol on diphenhydramine hydrochloride drug in aqueous solution at different temperatures: Volumetric and acoustic approach**, *J. Chem. Thermodyn.*, 87, (2015), pp. 78.
13. H. Kumar, K. Kaur **Interaction of antibacterial drug ampicillin with glycine and its dipeptides analyzed by volumetric and acoustic methods at different temperatures**, *Thermochim. Acta*, 551, (2013), pp. 40.
14. R. B. Sawant **Viscometric and speed of sound studies of binary mixtures of tert-butyl acetate with fluorobenzene, chlorobenzene and bromobenzene at (298.15 and 308.15) K and at atmospheric pressure 0.087 MPa**, *J. Solut. Chem.*, 47, (2018), pp. 795.
15. P. Bernal, L. Brown **Molar volumes and isentropic compressions of cyclic ethers in aqueous solutions from 288.15 to 313.15 K at atmospheric pressure**, *J. Solut. Chem.*, 47, (2018), pp. 408.
16. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji **Expanding the limits of computational chemistry**, Gaussian 09, Revision C.01, Gaussian 09, Revision C.01, Gaussian Inc.: Wallingford, CT, USA, (2009).
17. A. D. Becke **Density-functional thermochemistry III the role of exact exchange**, *J. Chem. Phys.*, 98, (1993), pp. 5648.
18. C. Lee, W. Yang, R. G. Parr **Development of the colle-salvetti correlation-energy formula into a functional of the electron density**, *Phys. Rev. B*, 37, (1988), pp. 785.

19. A. Soto, A. Arce, M. K. Khoshkbarchi **Thermodynamics of diglycine and triglycine in aqueous NaCl solutions: apparent molar volume, isentropic compressibility, and refractive index**, J. Solut. Chem., 33, (2004), pp. 11.
20. S. Chauhan, K. Singh, K. Kumar, S. C. Neelakantan, G. Kumar **Drug– amino acid interactions in aqueous medium: volumetric, compressibility, and viscometric studies**, J. Chem. Eng. Data, 61, (2016), pp. 788.