



A preliminary survey for hard ticks (Acari: Ixodidae) diversity on the livestock in Akola District, Maharashtra. (India)

Pooja S. Thakur, Nagrale S. M* and Raja. I. A.

Shri Shivaji college of Art Commerce and Science, Akola

*RLT Science college, Akola.

Abstract

Hard ticks (Acari: Ixodidae) are ectoparasitic feeders on blood, they carry and transmit a wide range of diseases of veterinary importance. Akola district of Maharashtra, from India, has a wide range of livestock which faces a problem of infestation from hard tick and other parasites. We surveyed and reported diversity of these hard ticks infesting major livestock from different localities in Akola district, in the year 2013. 255 livestock, consisting of 71 cattle, 45 buffaloes, 86 goats, and 53 sheep of all 7 talukas of the district were searched and total 590 adult ticks belonging to six species were sampled and were identified at species levels. All the ticks, belonging to three genera: Hyalomma (Bont-legged ticks), Rhipicephalus (Brown ticks) and Haemaphysalis. The species of ticks documented from Akola district were, *Hyalomma anatolicum anatolicum*, *Hyalomma marginatum isaaci*, *Hyalomma bussan*, *Rhipicephalus boophilus micropilus*, *Rhipicephalus haemaphysaloides*, and *Haemaphysalis hispanosa*.

Key words: Acari, Ixodidae, Akola, Diversity, Livestock

Introduction:

All ticks are ectoparasitic obligatory feeders of blood. They are sanguivorous on vertebrate host: mammals' birds and reptiles. Along with spiders, Ticks included in class Arachnida in phylum Arthropoda (Soulsby, 1982). They belong to subclass Acari which consists of many orders of mites and one order of tick, the Ixodida. Though there are three families of ticks but two are important for domestic animals (Sonenshine, 2014). Family Argasidae contains the important genera Argas, Ornithodoros, and Otobius. These genera are known as soft ticks. The family Ixodidae comprised of 13 genera and approximately 650 species (Sonenshine, 2014). It contains the important genera Amblyomma, Dermacentor, Haemaphysalis, Hyalomma, Ixodes, Margaropus, and Rhipicephalus. These genera are known as hard ticks because their outer surfaces have hard plates. Also, the important boophilid ticks, formerly of the genus Boophilus, are now classified as a subgenus within the genus Rhipicephalus (Tailor, 2007). Within these genera, more than 100 species are reported to infest domestic animals. Some of these species also feed on humans (Tailor, 2007). Boophilid ticks, a subgenus within Rhipicephalus ticks, commonly known as cattle ticks or blue ticks, have a highly characteristic morphology and one-host lifecycle causing direct parasitic losses and by transmission of microbe. Hyalomma genus contains many species of hard ticks important to domestic animals in hot dry regions in India, Africa, Mediterranean basin, and the Middle East. Genus Haemaphysalis found in Asia, Europe, Africa, and Australia. They affect cattle, sheep, goats, horses, camels, dogs, cats, and humans, as well as large range of wild mammals, birds and reptiles. Being obligate parasites ticks infest almost all sorts of livestock affecting greatly their health by blood sucking and transmitting diseases like babesiosis, theileriosis, anaplasmosis etc (Jongejan and Uilenberg,

2004). An estimated annual loss of US\$ 500000 is reported alone from low quality hides and skin in eastern Ethiopia (Desalegn *et al.*, 2015) on account of tick infestation. Since beginning of 20th century few experts studied ticks from different parts of India (Sharif, 1928 : Sen, 1938 : Dhanda and Rao 1964: Varma and Mahadevan 1970 : Kumar, *et al.*, 2002 : Patel *et al.*, 2013 : Singh and Rath 2013) and reported 109 species. Out of these 28 species of ticks reported to documented from Maharashtra by Sherif (1928): Sen, (1938) : Srivastava and Wattal, 1973: Mishra, *et al.*, (1977) : Raote, (1983) : Shahardar, (1988) : Varghese and Dhanda 1995).

Ticks as an obligatory ectoparasite of livestock, have high medical and veterinary importance, distributed all over the world. But their occurrence and prevalence differ greatly from region to region depending on various factors. In the present study a preliminary survey of tick is presented on the diversity and prevalence of ticks' species infesting major livestock in Akola district of Vidarbha region from Maharashtra. Study like this is locally important the studied district is a home for about 134481 goats, 49455 buffaloes, and 15467 exotic cross, and 269343 indigenous breed of cattle, (Govt livestock census report, 2012).

Materials and Methods.

Study Area:

The district of Akola lies in the western parts of the Vidarbha region of Maharashtra State and is surrounded by Amravati district in the north and north-east, Yeotmal in the south-east, Washim in the south and Buldhana in the west. The district lies between 19 ° - 51' and 21 ° - 16' latitude and 76 ° - 38' and 77 ° - 44' longitude. Though a large section of rural population depends on agriculture many of them also depend on livestock as additional livelihood. Thus these livestock plays an important role in the people's economy in the district.

Methodology:

This study was carried out, in the year 2013 in the district of Akola. The study commenced with a survey of all the talukas (Akola, Akot, Balapur, Barahi-Takli, Murtijapur, Patur, and Telhara,) of the district and information about the owners and their livestock, was collected. With the permission of the owner, frequent visits were made to some randomly selected localities/herds and all animals present were physically examined for their health status and tick attacks if any. Number of ticks infesting the host was counted and was pulled carefully out of the skin, put separately in glass vials containing 70% alcohol and 5% glycerol with proper labeling. Local Veterinary health centers were also visited and ticks were collected from different livestock. A total number of 594 hard ticks were collected from total 255 livestock, including goats, sheep, cattle and buffaloes. All are brought to Research laboratory, of Department of Zoology, Shri Shivaji College Akola. Identification was made under a stereomicroscope on the basis of shape of scutum, leg color, body, coxae one and ventral plates, with the help of the available standard keys. The identified ticks were photographed and the specimens are deposited with the departmental museum.

Results and Discussion.

The present study reported a survey of diversity of the tick infesting major livestock in Akola district of Vidarbha region in Maharashtra, India, in the year 2013. A total of 255 livestock, consisting of 71 cattle, 45 buffaloes, 86 goats, and 53 sheep were searched from all the seven talukas of the district and 143 were found infested with ticks. As depicted in the table.1 total 594 adult ticks, belonging to 6 species from three genera of Ixodid ticks were collected. The preferred predilection sites, of attachment of ticks to host were ears, axilla, groin, genital areas, perineum, udder, anal and peri-anal, abdomen, neck, shoulder, belly, flanks, back, upper legs, dewlap and hindquarters.

In our survey we reported prevalence of three genera: *Hyalomma* (Bont-legged ticks), *Rhipicephalus* (Brown ticks) and *Haemaphysalis*, it is in accordance with that of Srivastava and Watal (1973) who reported similar genera of ticks from, eastern Vidarbha region of Maharashtra. In our survey these genera are represented by six species of tick (Table.1: Photo Plate.1), which are *Hyalomma anatolicum anatolicum*, *Hyalomma marginatum issaaci*, *Hyalomma hussaini*, *Rhipicephalus boophilus micropilus*, *Rhipicephalus haemaphysaloides*, and *Haemaphysalis bispinosa*. Prevalence of these species from different parts of Maharashtra is also reported by Varghese and Dhanda (1995) in their documented 21 species of ticks on domestic animals.

Out of 71 cattle searched, 59 were positive for Ixodid ticks. In all, 6 Ixodid species, comprising 414 adult ticks, were collected from positive cattle.

Among 45 buffaloes we found 29 positive for ticks attack. Total 85 adult ticks were collected from 29 buffaloes. All studied six species of tick are found to infest both cattle and buffaloes. In cattle the order of prevalence was *Rhipicephalus boophilus micropilus*, *Hyalomma a. anatolicum*, *Haemaphysalis bispinosa* and *Rhipicephalus haemaphysaloides*, *Hyalomma hussaini* and *Hyalomma marginatum issaaci*. In buffaloes the order of abundance was *Rhipicephalus boophilus micropilus*, *Hyalomma a. anatolicum*, *Haemaphysalis bispinosa* and *Rhipicephalus haemaphysaloides*, *Hyalomma marginatum issaaci* and *Hyalomma hussaini* (Table.1). In our report the infestation rate of cattle was (83.04 %) and that of buffaloes was (64.44 %). However in an earlier report, from Maharashtra Varghese and Dhanda in 1995 found that infestation rate for buffalo in Maharashtra was 42.6% and that of cattle was 75.5%. The lower rate of infestation on buffaloes in our report might be due to more care and regular cleanliness by the owners. In contrast to our reports, Chhillar et al., (2014) reported lower infestation of both cattle and buffaloes, (cattle 55% and buffaloes 42%) from Haryana which is might be due to climatic conditions, in the northern India. Even after the treatment of acaricides, Jawale et al., (2012) reported that in Nashik district of Maharashtra 62% cattle, shows tick infestation.

As depicted in Table.1, total, 53 sheep were searched in Akola district, and out of which 27 were positive for ticks and out of 86 searched goats, 28 were positive. In all 46 and 45 adults, ticks, belonging to four species of ticks, were found on sheep and goats respectively. Out of them, *Haemaphysalis bispinosa*, and *Rhipicephalus haemaphysaloides* are most dominant ticks on both goats and sheep, while the prevalence of *Hyalomma a. anatolicum* and *Hyalomma hussaini* is quite lower. The findings are in accordance with that of Diyes and Rajakaruna, (2015) who too reported that the dominant genera ticks infesting the small livestock like goats are, *Haemaphysalis*, and *Rhipicephalus* from the nearby island of Srilanka and it was the, *Haemaphysalis bispinosa*, which second most dominant species attacking goats. Srinadarajan

et al., (2011) shows that *H. bispinosa* parasitize more goats like mammals than any other domesticated animal in the state of Tamilnadu, and that it is widely distributed. In the present study too *H. bispinosa* was the most prevalent species on the goats of Akola district. We reported that the tick infestation of sheep is 50.94 % and that of goats, is 32.55 % which is much comparable with that of Varghese and Dhanda (1995) who too reported 53.90 % infestation of Sheep and 36.20 % infestation in goats, from the same state of Maharashtra. Here we also reported that all the studied livestock, infested by all the identified species of ticks in Akola district, which is due to the fact that, in many areas cattle, buffaloes, goats and sheep, are usually kept together in vicinity of each other.

even they use common grazing grounds. So the ingesting ticks adapted for to multi host parasitism.

Being obligatory haematophagous ectoparasite of veterinary importance, result in the present survey of tick from Akola district becomes increasingly significant and will definitely prove helpful in promoting control measures of tick and tick-borne disease among the local livestock.

References:

Chhillar Shivani, Chhilar, J. S and , Kaur Harpreet. (2014). Investigations on Some Hard Ticks (Acari: Ixodidae) Infesting Domestic Buffalo and Cattle from Haryana, India. *Journal of Entomology and Zoology Studies*; 2 (4): 99-104

Desalegn T, Fikru A and Kasaye S (2015). Survey of Tick Infestation in Domestic Ruminants of Haramaya District, Eastern Hararghe, Ethiopia. *Journal of Bacteriology and Parasitology* 6, 246-49.

Dhanda V. and Rao T. R.(1964). "A report on collection of Ixodid ticks made in North East Fortier Agency India," *Indian Journal of Medical Research*, vol. 52, pp. 1139-1153,

Diyas, G. C. P and Rajakaruna. R. S. (2015) Diversity and distribution of tick species infesting goats with two new host records from Sri Lanka. *J.Natn.Sci.Foundation Sri Lanka* 2015 43 (3):225-234

Jawale C. S., Vinchurkar A. S., Dama L. B. and Dama S.B. 2012.Prevalence of ixodid ticks in post acaricide treated cattle and buffaloes at Sinner District Nashik (M.S) India. *Trends in parasitological research*. Vol. 1 No. 1. p.20-24.

Kaushal Kumar, Balakrishnan N., and Sharma Abhay Kumar. Kiszewski. 2014. Studies on the Vertical Distribution of Ticks of Domestic Animals and Their Public Health Importance in Nilgiri Hills and Adjoining Areas of Tamil Nadu State (India). *Hindawi Publishing Corporation International Journal of Zoology* Volume 2014, Article ID 359812, 6 pages .

Kiszewski, A. E., Matuschka, F and Spielman, A. (2001). Mating strategies and spermiogenesis in Ixodid ticks. *Annual Review of Entomology*, 46(1), 167-182.

Kumar, K., Balakrishnan N., Katyal R., and Gill, K. S. (2002). Prevalence of Ixodid ticks in Nilgiri District of Tamil Nadu State (India)." *Journal of Communicable Diseases*, vol. 34, no. 2, pp. 124-127,.

Patel G., Shanker D., Jaiswal A. K., Sudan V., and Verma S. K., (2013). "Prevalence and seasonal variation in ixodid ticks on cattle of Mathura district, Uttar Pradesh," *Journal of Parasitic Diseases*, vol. 37, no. 2, pp. 173-176,

Rao, Y. V. (1983).Incidence of parasitic infestation on organized cattle farms in Bombay region. M.V.Sc. thesis submitted

to Konkan Krishi Vidyapeeth, Dapoli, Ratnagiri (Not seen)

A. K. Sanyal and S. K. De. 2004. *Acari: Metastigmata (Ticks)* Zool. Surv. India State Fauna Series 8: Fauna of Gujarat 11-18.

Sen, P., (1938). A check and host-list of Ixodoidea (Ticks) occurring in India. *Indian J. Vet. Sci. Anim. Husbandry*, 8: 133-149.

Sharif, M., (1928). A revision of the Indian Ixodidae with special reference to the collections in the Indian museum. *Rec. Indian Mus.*, 30: 217-344.

Shahardar, F. A. Studies on ixodid ticks of bovines in Konkan region of Maharashtra state. M.V.Sc. thesis submitted to Konkan Krishi Vidyapeeth, Dapoli, Ratnagiri (1988). (Not seen)

Singh N. K. and Rath S. S., (2013). "Epidemiology of ixodid ticks in cattle population of various agro-climatic zones of Punjab, India," *Asian Pacific Journal of Tropical Medicine*, vol. 6, no. 12, pp. 947-951,

Soulsby, E. J .L.(1982): *Helminths, Arthropods and Protozoa of Domesticated Animals*,7th edition the ELBS and Baillieve, Tindal cass LTD, London, pp.766.

Soundararajan C., Latha B. R. and Pandian S. S. (2014). Prevalence of tick infestation in goats under different system of management. *International Journal of Agricultural Science and Veterinary Medicine* 2: 2Sonenshine, D.E. (2014). *Biology of ticks* (vols 1&2). New York: Oxford University Press. ISBN 978 0-19-974405-3

Srivastava, S. P. and Wattal B. L. (1973). Studies on the distribution of ectoparasites and their vertebrate hosts in urban, semi urban (peripheral) and sylvatic zone of Nagpur city and neighbouring areas (Maharashtra, India). - *Indian J. Ent.*, 35 (4): 306-324.

Taylor, M. A. (2007). *Veterinary parasitology*. Oxford: Blackwell Publishing. ISBN 978-1-4051-1964-1.

Varma R. N. and Mahadevan B., (1970). "Ixodid ticks collected in the Eastern Himalayas and their potential disease relationships," *The Indian Journal of Medical Science*, vol. 58, no. 6, pp. 693-706

Varghese G. And V. Dhanda . (1997). *Ixodid Ticks of Maharashtra State*. *India. Acarologia*, t. XXXVI, fasc. 4, (1995).

Walker A. R., Bouattour A., Camicas J. L., A. Estrada-Pena., Hornk I. G., Lutf A. A.,Pegram R. O. and Preston P. M. (2014). *Ticks of Domestic Animals in Africa: a Guide to Identification of Species*. Bioscience Reports, Edinburgh Scotland, U.K. www.biosciencereports.pwp.blueyonder.co.uk